

Klean-Prep[®] for Capsule Endoscopy

Bowel Preparation Instructions

This leaflet gives you instructions on how and when to take your bowel preparation medication. It is important that you follow these instructions carefully. If your bowel preparation is not clear then a diagnosis may not be made and you may need to undergo the procedure again.

7 days before Colonoscopy

- Stop iron tablets (Ferrous Sulphate) some multivitamins or supplements may contain small amount of iron
- Stop any food containing bran (oat bran, wheat bran, rice bran etc.)

4 days before Colonoscopy

- Stop taking constipating agents such as Lomotil, Imodium/Loperamide
- Stop taking pain relief medication containing Codeine Phosphate (Co-codamol[®]/Co-dydramol[®] etc.)
- Please continue taking all your other prescribed medication

3 days before Colonoscopy

- Try drinking at least 2 litres of clear fluids (8-10 glasses) per day. List of clear fluids allowed as below.
- You can continue to eat your normal meals but please avoid foods that are high in fibre. (See list below)

	ALLOWED (✓)	AVOID (X)
Cereals	Rice Crispies, Frosties, Sugar Puffs, Coco Puffs, Cornflakes	All wheat based cereals like Shredded Wheat, Weetabix, porridge, Fruit and Fibre [®] , Muesli, granola, wheat bran and all bran
Pasta	white pasta	wholemeal pasta
Rice	white rice	brown rice
Bread	white bread	wholemeal, granary, soft grain, oat bread
Meat	chicken, turkey, fish, eggs	All red meat
Fruits and Vegetables	Bakes, boiled, mashed potatoes WITHOUT skin	All fruits and vegetables
Dairy	Butter, margarine, cheese	
Puddings /cakes	Mousse, jelly (not red), sponge cakes, madeira cakes. Rich tea biscuits, wafer biscuits	Digestive, oat biscuits, Ryvita or anything containing wholemeal flour, dried fruits and nut

	ALLOWED (✓)	AVOID (X)
Soups	Sieved soup, clear broth, stocks Marmite® /Bovril® /OXO® mixed with hot water	Chunky vegetable or lentil soups
Drinks	clear fluids include squash drinks, coke, lemonade, tonic/soda, Lucozade, jelly, black tea or coffee	Milk, fruit juices with pulp, tomato juice
Preserves and Sweets	Sugar, jam, marmalade, honey, maple syrup, lemon curd	Jams or marmalade with peels and seeds, chocolate bars containing dried fruits and nuts
Others	Soya, tofu	Nuts, dried fruits, Quorn, hummus

1 day before your appointment

Please follow the instructions below and tick appropriate box once you have started the regimen.


Time	Afternoon Appointment	Please tick
7am	Good breakfast (from the list provided) then nothing solid afterwards	
2 pm	Take one sachet of Klean-Prep and drink a glass every 15 minutes. (Please see instructions below)	
5pm	Take one sachet of Klean-Prep and drink a glass every 15 minutes. (Please see instructions at the back)	

On the day of your procedure

Do **NOT** eat or drink anything, except essential medication with a sip of water.

Instructions on how to prepare Klean – Prep


This solution taste better when chilled so you may wish to prepare this in advance. Dissolve the first sachet in a litre of warm water to dissolve the contents. Leave it to cool and drink one glass every 10 - 15 minutes. You may find it easier to drink with a straw and if the mixture is chilled. You could also add a dash of fruit squash, sugar or honey to make it taste nicer. If you feel sick while drinking the preparation, stop temporarily. Keep moving about and you should feel better.



It is advisable to stay near the toilet once you have started taking your bowel preparation. You should expect frequent bowel movement and diarrhoea within one to three hours from the first dose of the medication, this will continue until you have taken the last dose of the medicine. You may want to apply some cream such as Zinc and Caster Oil in your bottom to prevent soreness.

Remember to keep drinking fluids especially once the diarrhoea has started to prevent you from getting dehydration. As a guide, you must drink at least two litres of clear fluids on the day that you are taking your bowel preparation.

Any questions regarding the above information, please ring us on 01932 722231, between the hours of 0800-1800 from Monday to Saturday.



Further Information

We endeavour to provide an excellent service at all times, but should you have any concerns please, in the first instance, raise these with the Matron, Senior Nurse or Manager on duty. If they cannot resolve your concern, please contact our Patient Experience Team on 01932 723553 or email asp-tr.patient.advice@nhs.net. If you remain concerned, the team can also advise upon how to make a formal complaint.

Author: Sister Jean Asare

Department: Endoscopy

Version: 1

Published: August 2018

Review: August 2020

We can provide interpreters for a variety of languages, information in larger print or other formats (e.g. audio) - please call us on 01932 723553.



اگر نیاز به ترجمہ دارید، لطفاً با شماره 01932 723553 تماس بگیرید۔
 ਜੇ ਤੁਹਾਨੂੰ ਡਰਾਈਵਿੰਗ ਦੀ ਲੋੜ ਹੈ ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਇਸ ਨੰਬਰ 'ਤੇ ਫੋਨ ਕਰੋ: 01932 723553
 اگر آپ اس کا اردو زبان میں ترجمہ چاہتے ہیں، تو براؤزر میں اس فون نمبر 01932 723553 پر رابطہ کریں
 Se precisa de uma tradução por favor contacte: 01932 723553

আপনার অনুবাদের দরকার হলে এখানে যোগাযোগ করুন : 01932 723553
 यदि आपको अनुवाद की ज़रूरत है तो कृपया इस नंबर पर फोन करें: 01932 723553
 Jeżeli chcemy, aby te informacje w innym języku,
 proszę zadzwonić 01932 723553

Ashford Hospital London Road, Ashford, Middlesex, TW15 3AA Tel: **01784 884488**
St. Peter's Hospital Guildford Road, Chertsey, Surrey, KT16 0PZ Tel: **01932 872000**

Website: www.ashfordstpeters.nhs.uk