



We can provide interpreters for a variety of languages, information in larger print or other formats (e.g. audio) - please call us on 01932 723553.

To use the Text Relay service, prefix all numbers with 18001.

اگر نیاز به ترجمہ دارید، لطفاً با شماره 01932 723553 تماس بگیرید.

ਜੇ ਤੁਹਾਨੂੰ ਤਰਜਮੇ ਦੀ ਲੋੜ ਹੈ ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਇਸ ਨੰਬਰ ਤੇ فون ਕਰੋ: 01932 723553

اگر آپ اس کا اردو زبان میں ترجمہ چاہتے ہیں، تو براہ کرم اس فون نمبر 01932 723553 پر رابطہ کریں

Se precisa de uma tradução por favor contacte: 01932 723553

আপনার অনুবাদের দরকার হলে এখানে যোগাযোগ করুন : 01932 723553

यदि आपको अनुवाद की ज़रूरत है तो कृपया इस नंबर पर फोन करें: 01932 723553

Jeżeli chcemy, aby te informacje w innym języku, proszę zadzwonić 01932 723553

Ashford Hospital
London Road
Ashford, Middlesex
TW15 3AA
Tel: **01784 884488**

St. Peter's Hospital
Guildford Road
Chertsey, Surrey
KT16 0PZ.
Tel: **01932 872000**

Website: www.ashfordstpeters.nhs.uk

Fistulogram, Fistuloplasty and Venoplasty Interventional Imaging Department



This leaflet tells you about having a fistulogram, fistuloplasty or venoplasty. It explains what is involved and what the possible risks are. It is not meant to replace informed discussion between you and your doctor, but can act as a starting point for such discussions. If you have any questions about the procedure please ask the doctor who has referred you or the department which is going to perform it.

What is a fistulogram?

This is an examination of the blood vessels that make up your fistula. A small needle is placed in your fistula and dye (contrast agent) is injected. This dye provides an image (like a map) of the blood vessels that would otherwise be invisible on X-ray. The interventional radiologist interprets the images.

What is a fistuloplasty or venoplasty or stent?

Occasionally, the blood vessels that make up a fistula can develop a narrowing which is diagnosed with a fistulogram. A special balloon called an angioplasty balloon is introduced into the area of narrowing inside the fistula. This balloon is inflated from outside the body momentarily and then deflated to improve the narrowing. A completion fistulogram is then carried out to check the result. If this is in the fistula it is called a fistuloplasty, if it is in a central vein it is called a venoplasty. Very rarely, a special metal tube called a stent is inserted into the fistula. This is permanent and keeps the narrowing open. Stents are generally

Finally

Some of your questions should have been answered by this leaflet, but remember that this is only a starting point for discussion about your treatment with the doctors looking after you. Make sure you are satisfied that you have received enough information about the procedure.

Reproduced with the kind permission of the British Society of Interventional Radiology (BSIR) 2011

Further Information

We endeavour to provide an excellent service at all times, but should you have any concerns please, in the first instance, raise these with the Matron, Senior Nurse or Manager on duty.

If they cannot resolve your concern, please contact our Patient Experience Team on 01932 723553 or email asp-tr.patient.advice@nhs.net. If you remain concerned, the team can also advise upon how to make a formal complaint.

Author: Interventional Team

Department: Imaging

Version: 3

Published: Mar 2018

Review: Mar 2020

groin as all the veins inside your body are connected and treatment is sometimes carried out via the groin vein, as this may be a safer option than directly through the fistula.

Will it hurt?

The local anaesthetic will sting initially, but this soon passes. Occasionally when the balloon is inflated during fistuloplasty, a dull ache may occur but this passes when the balloon is deflated.

How long will it take?

Every patient is different, and it is not always easy to predict; however, expect to be in the radiology department for around 15–20 minutes for a fistulogram and fistuloplasty or venoplasty can take up to an hour.

What happens afterwards?

Light pressure is applied for a few minutes to the region where the needle/plastic tube was placed to prevent bleeding. You can usually go home 30 minutes after a fistulogram, but will need to stay in hospital for a few hours following fistuloplasty or venoplasty. A follow-up fistulogram will usually be required after one month. You will receive this appointment in the post.

only used if the balloon does not improve the narrowing satisfactorily or if there is a complication.

Why do you need a fistulogram?

Your doctor feels that there may be a problem with your dialysis fistula (or graft). This test is the best way to diagnose the problem associated with your fistula. You may already have had a Doppler ultrasound scan to assess the flow in your fistula to help decide how to approach this problem.

What are the risks of a fistulogram?

Fistulograms are very safe procedures but there are some risks and complications that can occur. Very occasionally, a small bruise can appear at the site of needle puncture. Less commonly, ongoing bleeding in this area leads to a short inpatient stay (once or twice in every hundred). Very rarely damage to the fistula can occur that may require further treatment by the interventional radiologist or a small operation. The risk of infection is very low.

What are the risks of a fistuloplasty/venoplasty?

Fistuloplasty/venoplasty, like a fistulogram, is very safe but occasionally complications do arise. A fistulogram is often performed before a fistuloplasty. There is a small risk of failure of treatment. Sometimes the narrowing in a fistula does not respond well to the fistuloplasty/venoplasty and requires a stent. The risk

of bleeding is slightly higher than for fistulogram (about three-in-100 chance). There is a small risk that the treatment may damage or even rupture the fistula/vein. If this were to happen, the fistula may fail and could not be used for dialysis. A small operation may be required at the time but more likely a line would be placed and a new fistula fashioned. When considering this risk, it is important to bear in mind that leaving a narrowing in a fistula or vein, without treatment it is likely your fistula would ultimately fail.

Who has made the decision?

Your doctors, the vascular access nurse and the interventional radiologist will have discussed your care and feel this is the most appropriate next step. However, you will also have the opportunity for your opinion to be considered and if, after discussion with your doctors, you no longer want the procedure, you can decide against it.

Are you required to make any special preparations?

Fistulograms are performed as an outpatient. However, if you may require a fistuloplasty or venoplasty, you may be admitted as a day case and asked not to eat or drink for four hours before the procedure. If you have any allergies or have previously had a reaction to the dye (contrast agent), you *must* tell the radiology staff before you have the test.

Who will you see?

A specially trained team led by an interventional radiologist within the radiology department. Interventional radiologists have special expertise in reading the images and using imaging to guide catheters and wires to aid diagnosis and treatment.

Where will the procedure take place?

In the angiography suite or theatre; this is usually located within the radiology department. This is similar to an operating theatre into which specialised X-ray equipment has been installed.

What happens during a fistulogram, fistuloplasty or venoplasty?

You will be asked to get undressed and put on a hospital gown. A small needle may be placed in your non-fistula arm in case you need an injection of a painkiller or light sedative. You will be asked to lie flat on your back. The skin over the area of your fistula will be swabbed with an antiseptic and you will be covered with sterile drapes. Local anaesthetic will be injected into the skin. A needle, often followed by a fine plastic tube, will then be placed in the fistula and dye injected. You will be asked to hold your breath for a few seconds while the images are taken. If you have a fistuloplasty or venoplasty, you will have monitoring devices placed on your chest and on your finger and if a sedative is given, you will be given oxygen via a face mask. Occasionally, it may be necessary to place a fine plastic tube in the vein in your